Start Time.	VENTILATORY MANAGEMENT		
Stop Time:	ENDOTRACHEAL INTUBATION		
Candidate's Name:	Date:		
Evaluator's Name:			
Note: If a candidate elects to initially ventilate the patient with a BVM attached to a reservoir and oxygen, full credit must be awarded for steps denoted by " ** " provided the first ventilation is delivered within the initial 30 seconds		Points Possible	Points Awarded
Takes of verbalizes body substance isolation precautions		1	
Opens the airway manually		1	
Elevates the patient's tongue and inserts a simple airway adjunct (oropharyngeal/nasopharyngeal airway)		1	
Note: The examiner must now int adjunct."	form the candidate "no gag reflex is present and the patient accep	ots the air	rway
** Ventilates the patient immediately using a BVM device unattached to oxygen		1	
** Hyperventilates the patient with room air		1	
Note: The examiner must now int	form the candidate that ventilation is being properly performed w	ithout dif	ficulty
Attaches the oxygen reservoir to the	e BVM	1	
Attaches the BVM to high flow oxygen (15 liter per minute)		1	
Ventilates the patient at the proper	volume and rate (800-1200 ml/breath and 10-20 breaths/minute)	1	
sounds are present and equal bil must now take over ventilation o		on. The e	
Directs assistant to hyper-oxygenate the patient Identifies/selects the proper equipment for endotracheal intubation		1	
	124 (PDIS) - 1 4 (1	
Checks equipment	Checks for cuff leaks Checks laryngoscope operation and bulb tightness	1	
	the OPA and move out of the way when the candidate is prepare	d to intu	bate the
patient. Positions the patient's head propert		1	
Inserts the laryngoscope blade into the patient's mouth while displacing the patient's tongue laterally		1	-
Elevates the patient's mandible with the laryngoscope		1	-
Introduces the endotracheal tube and advances the tube to the proper depth		1	-
Inflates the cuff to the proper pressure		-	+
		1	-
Disconnects the syringe from the cuff inlet port		1	-
Directs assistant to ventilate the patient Confirms proper placement of the endotracheal tube by auscultation bilaterally and over the epigastrium		1	-
	you had proper placement, what would you expect to hear?"	1	
Secures the endotracheal tube (ma		1 1	Т
	***************************************	-	+
Did not initiate ventilation wit	Total: y substance isolation precautions when necessary hin 30 seconds after applying gloves or interrupts ventilations for great	21 ter than 30	0
seconds at any time Did not voice or provide high Did not ventilate the patient a Did not provide adequate vol Did not hyper-oxygenate the Did not successfully intubate Used the patient's teeth as a Did not assure proper tube p The stylette (if used) extends	oxygen concentrations (15 liter/minute or greater) at a rate of at least 10 breaths per minute ume per breath (maximum of 2 errors per minute permissible) patient prior to intubation the patient within 3 attempts		